

AMS Joint Meeting
San Francisco, 2010

First coefficient of the
Conway polynomial of virtual
links

Sergei Chmutov

Ohio State University, Mansfield

January 14, 2010

9:30 - 9:50

Joint work with
Z. Cheng, T. Dokos, J. Lindquist ^U

I. Classical links

L link with m components

$$\nabla(L) = c_0 + c_1 z + c_2 z^2 + \dots$$

the Conway polynomial

I.1 Theorem

(F. Hosokawa '58, R. Hartley '83,
J. Hoste '85)

$$c_0 = c_1 = \dots = c_{m-2} = 0$$

$$c_{m-1} = \det \Lambda^{(p)}$$


$$\Lambda = (\lambda_{ij}), \quad \lambda_{ij} = \begin{cases} -lk_{ij}(L) & \text{if } i \neq j \\ \sum_{k \neq i} lk_{ik}(L) & \text{if } i = j \end{cases}$$

$$\Lambda^{(p)} = \left(\begin{array}{c} \dots \\ \dots \\ \text{X} \\ \dots \\ \dots \end{array} \right) \text{ p-th}$$

p-th

Matrix-Tree Theorem

(2)



$$\det \Delta^{(P)} = \sum_{\substack{T \\ \text{spanning} \\ \text{tree}}} \prod_{e \in T} w_e$$

I.2

algebraically split links
 $l_{ij} = lk_{ij}(L) = 0$
for all i, j .

Theorem (L. Traldi '84, J. Levine '97)

$$C_{m-1} = C_m = \dots = C_{2m-3} = 0$$

$$C_{2m-2} = \det M^{(P)}$$

$$M = (m_{ij}), \quad m_{ij} = \sum_k M_{ijk}(L)$$

triple Milnor numbers

$$+M_{ijk}(L) = -M_{jik}(L) = +M_{jki}(L)$$

G. Masbaum, A. Vaintrob '02

(3)

Pfaffian Matrix-Tree Theorem

II Virtual links

II.1 Linking numbers $l_{i/j} \neq l_{j/i}$

$$l_{i/j} := \sum_{\substack{\nearrow \\ \searrow \\ i \quad j}} \varepsilon_{\nearrow/\searrow}, \quad \begin{array}{c} \nearrow \\ \oplus \\ \searrow \end{array} \quad \begin{array}{c} \nearrow \\ \ominus \\ \searrow \end{array}$$

Conway polynomial

(—, M. Khoury, A. Rossi '09)


$$\nabla_{\text{asc}}(L_*) = \sum_S \left(\prod_{\nearrow/\searrow \in S} \varepsilon_{\nearrow/\searrow} \right) \cdot \mathbb{Z}^{|S|}$$

ascending
one-component

- Smooth the crossings from S according to the orientation.
- S is one-component if the link obtained is a knot.
- S is ascending if at the first approach to each crossing of S we jump down to smooth it.


Example.

$l_{1/2} = -1, l_{2/1} = 0,$
 $l_{1/3} = 0, l_{3/1} = -1,$
 $l_{2/3} = -1, l_{3/2} = 0$




$\Delta_{asc}(L) = \mathbb{Z}^2$

$S = \{a\}$



not one-component
but ascending

$S = \{a, b\}$



one-component
ascending


Theorem

(5)

$$C_{m-1} = \det \Lambda^{(1)}$$

$$\Lambda = (\lambda_{ij}), \quad \lambda_{ij} = \begin{cases} -e_{j/i}, & \text{if } i \neq j \\ \sum_{k \neq i} e_{k/i}, & \text{if } i = j \end{cases}$$

Oriented matrix-tree Theorem



$$\det \Lambda^{(1)} = \sum_T \prod_{e \in T} w_e$$


oriented
spanning tree
growing from
vertex 1

Example $m=3$

(6)

$$\Delta = \begin{pmatrix} l_{3/1} + l_{2/1} & -l_{2/1} & -l_{3/1} \\ -l_{1/2} & l_{1/2} + l_{3/2} & -l_{3/2} \\ -l_{1/3} & -l_{2/3} & l_{1/3} + l_{2/3} \end{pmatrix}$$
$$= \begin{pmatrix} -1 & 0 & 1 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}$$

$$\det \Delta^{(1)} = l_{1/2} l_{1/3} + l_{1/2} l_{2/3} + l_{3/2} l_{1/3}$$



II.2

Work in progress