Correlation between pole location and asymptotic behavior for Painlevé I solutions

Ovidiu Costin
Department of Mathematics
The University of Chicago

Abstract

We extend the technique of asymptotic series matching to exponential asymptotics expansions (transseries) and show that the extension provides a method of finding singularities of solutions of nonlinear differential equations, using asymptotic information. This transasymptotic matching method is applied to Painlevé's first equation. The solutions of P1 that are bounded in some direction towards infinity can be expressed as series of functions, obtained by generalized Borel summation of formal transseries solutions; the series converge in a neighborhood of infinity. We prove (under certain restrictions) that the boundary of the region of convergence contains actual poles of the associated solution. As a consequence, the position of these exterior poles is derived from asymptotic data. In particular, we prove that the location of the outermost pole $x_p(C)$ on $\mathbb{R}^+$ of a solution is monotonic in a parameter C describing its asymptotics on antistokes lines[1], and obtain rigorous bounds for $x_p(C)$. We also derive the behavior of $x_p(C)$ for large $C \in \mathbb{C}$. The appendix gives a detailed classical proof that the only singularities of solutions of P1 are poles.

1 Introduction and notations

We consider the solutions of Painlevé's first equation, P1

$$y'' = 6y^2 + z \tag{1.1}$$

and study the connection between their asymptotic behavior and the location of their poles.

The analysis and proofs in the present paper are mostly self-contained, but the technique used, as well as the nature of results obtained, are much easier to interpret from the more general perspective of exponential asymptotics. We start by briefly presenting some concepts and results in this area which are relevant to the problem under discussion.

Jean Écalle has introduced and developed the theory of analyzable functions and tools of generalized resummation of *transseries* (generalized exponential-power series) as a natural and very comprehensive generalization of the theory of analytic functions [8, 11, 12]. The grand picture is the construction of a class of functions, closed under all imaginable algebraic

and analytic operations, which are into an isomorphic correspondence with the space of transseries. In a sense, the space of transseries represents the formal closure of power series under all operations. A wide class of analytic problems admit formal solutions in the space of transseries (obtained in a more or less straightforward way). Transseries solutions contain ample analytic information and in many cases (for instance generic linear or nonlinear differential systems [16]) the actual solutions of the problem can be reconstructed from their transseries by modern resummation methods. Unlike the isomonodromic deformation approach (see [2, 10, 5, 9, 4]) the method introduced in this paper is suitable to be used in nonintegrable systems as well.

1.1 Notes on asymptotics beyond all orders for P1

Proofs of the various technical statements in this section can be found in [16], in the more general context of analytic rank-one differential systems.

The change of variables

$$\pm i\sqrt{6z^{-1}}y(z) = -1 + \frac{4}{25x^2} - h(x);\ 30x = (-24z)^{5/4} \tag{1.2}$$

brings (1.1) to the (normal) form

$$h'' + \frac{1}{x}h' - h - \frac{1}{2}h^2 - \frac{392}{625x^4} = 0 \tag{1.3}$$

There is a one-parameter family of solutions of (1.3) that tend to zero as $x \to +\infty$, $x \in \mathbb{R}$, and they have a common asymptotic series (cf. also § 3):

$$h(x) \sim \tilde{h}_0(x) = \sum_{k=4}^{\infty} c_k x^{-k} \tag{1.4}$$

The classical expansion (1.4) does not distinguish different solutions in this family. However there exists a one complex parameter family of transseries solutions, of which $\tilde{h}_0$ is only a special instance:

$$\tilde{h}(x, C) = \tilde{h}_0(x) + \sum_{k=1}^{\infty} C^k x^{-k/2} e^{-kx} \tilde{h}_k(x) \quad (C \in \mathbb{C}) \tag{1.5}$$

In (1.5), $\tilde{h}_k$ are formal power series, $\tilde{h}_k = \sum_{m=0}^{\infty} c_{km} x^{-m}$, that can be computed algorithmically by substituting (1.5) into (1.3) and identifying the coefficients. With the choice $c_{10} = 1$, which is possible for (1.3), the series $\tilde{h}_k$ do not depend on C. Every $\tilde{h}_k$ is factorially divergent. There is nevertheless a very close connection between (1.5) and the family of actual solutions of (1.3) which decrease on $\mathbb{R}$ for $x \to +\infty$: any such solution can be reconstructed by generalized Borel resummation of $\tilde{h}(x, C)$ for some C. Denote by $\mathcal{LB}$ the generalized Borel resummation operator (as defined in [16]). There exists $x_0 > 0$ (independent of k) so that the functions $h_k = \mathcal{LB}\tilde{h}_k$ are analytic in the half plane $H_{x_0} = \{x : \Re(x) > x_0\}$. In addition, $h_k \sim \tilde{h}_k$ as $x \to e^{i\theta}\infty$ in H_{x_0}, $\theta \in (-\pi/2, \pi/2)$. For any $C \in \mathbb{C}$, the function series

$$h(x; C) = \mathcal{LB}\tilde{h}_0(x) + \sum_{k=1}^{\infty} C^k x^{-k/2} e^{-kx} \mathcal{LB}\tilde{h}_k(x) \tag{1.6}$$

converges for $\Re(x) > x_1 = x_1(C)$ to a solution of (1.3); conversely, any solution h which vanishes for real $x \to +\infty$ is necessarily of the form (1.6) for a unique C. $\mathcal{LB}$ commutes with the algebraic operations, complex conjugation (thus a series with real coefficients sums to a real function), with differentiation, and has good continuity properties. Along directions in $\mathbb{C}$ other than $\mathbb{R}^+$ similar results hold.

In addition, the constant C in (1.6) is related by a simple formula to the large $|x|$ behavior of $h(x; C)$ on the antistokes lines $\pm i\mathbb{R}^+$ (cf. (2.8)). We show that for any $C > 0$, the boundary of the domain where the series (1.6) converges contains an actual pole of $h(x; C)$, much as the boundary of the convergence disk of the Taylor series of an analytic function must contain a singularity.

1.2 Heuristic discussion of transasymptotic matching

The value of C is related to the position of the outermost singularities of solutions of P1. The following heuristic argument illustrates the reason.

For simplicity, we assume that C is large. When x is very large, the exponentials in (1.5) are smaller than the leading term in $\tilde{h}_0$, and the behavior of $h(x; C)$ is roughly $c_{04} x^{-4}$.

For smaller values of x, such that $1 \gg |Ce^{-x}x^{-1/2}| \gg |x|^{-1}$, the relative magnitude of the terms of (1.5) changes: the largest term becomes $Ce^{-x}x^{-1/2}$ (instead of $c_{04}x^{-4}$), the next order of magnitude is the term $C^2e^{-2x}x^{-1}c_{20}$ (not $Ce^{-x}x^{-3/2}c_{11}$) and so on. In this intermediate regime,

the terms of (1.5) need to be reshuffled accordingly, and $h(x;C)$ is represented by

$$(1.7)\quad \tilde{h}(x;C) = \sum_{m=0}^{\infty} \frac{1}{x^m} \left(\sum_{k=0}^{\infty} \frac{c_{km}}{(e^x x^{1/2}/C)^k} \right) := \sum_{m=0}^{\infty} \frac{G_m(e^x x^{1/2}/C)}{x^m}$$

It turns out that, for x not too small, the inner sums converge (while, obviously, the expansion $\tilde{h}$ as a whole is still divergent), and $G_m(t)$ are analytic. In fact, for (1.3), $G_m(t)$ are rational functions and G_0 has a particularly simple expression, $G_0(t) = t(t-1/12)^{-2}$ (see the proof of Lemma 7). Moreover, for any $m \geq 0$ the function $G_m(t)$ has only one pole in the right half plane, at $t = 1/12$.

The representation (1.7) is normally accurate as long as no Stokes phenomena occur and the following asymptoticity condition holds

$$(1.8)\qquad \left| G_{m+1}(e^x x^{1/2}/C) x^{-m-1} \right| \ll \left| G_m(e^x x^{1/2}/C) x^{-m} \right|$$

A formal calculation shows that (1.8) holds for x in a region $\mathcal{R}$ that starts well to the *left* of the array of poles of $g(x) = G_0(e^x x^{1/2}/C)$, located at x_N, where

$$(1.9)\qquad x_N + \frac{1}{2} \ln x_N = \ln(C/12) + 2N\pi i;\ N \in \mathbb{Z}$$

and extends all the way (small neighborhoods of the poles themselves excluded) to $\Re(x) \to \infty$. The region $\mathcal{R}$ is thus substantially wider than the region where (1.4) is valid.

Accepting for the moment that (1.7) is the asymptotic expansion of some solution h in $\mathcal{R}$, (1.7) can be used to estimate $\oint(\int h)$ on a circuit around a pole x_N. The calculation gives $\oint(\int h) = -12\pi i + O(1/\ln C)$ for large $|C|$, showing that h is singular at some point $x_p(C;N) = x_N + o(1)$. $x_p(C;N)$ can be estimated to all orders in $1/\ln C$ from (1.7).

On the other hand, if x is large, and away from x_N, $h(x;C)$ must be bounded (by (1.7)). The location of the poles of h in $\mathcal{R}$ should thus be $x_p(C;N) = x_N + o(1)$.

This method gives information not only about the location, but also about the nature of the singularities of $h(x;C)$ in $\mathcal{R}$. A straightforward calculation based on the Cauchy integral argument explained above indicates, to any order in $1/\ln C$, that $x_p(C;N)$ are double poles.

The fact that P1 is an integrable equation plays no role, except that G_m can be computed in closed form.

We note that in the matching process *all the terms* of the transseries (1.5) are necessary, so that the matching is *trans*asymptotic. The procedure makes the connection between the behavior of h in the pole-free region near infinity, and its singularities for large x in $\mathbb{C}$.

In the next sections these conclusions are made rigorous, and further results are proved on the connection between C and the position of $x_p(C;0)$, for C not necessarily large. The proofs yield global estimates for (1.6) and (1.7).

We can also interpret (1.7) from the perspective of multiscale analysis. Apparently an expansion of the form $\sum c_m(x)x^{-m}$ (where the coefficient of x^{-m} may depend on x) is subject to substantial arbitrariness. Not so, however, if $c_m(x)$ are required to be sufficiently regular and the expansion to be valid over a wide enough range of x. This is a crucial point in multiscale analysis. We note that (1.5) determines the asymptotic scales x^{-1} and $Ce^{-x}x^{1/2}$. Once the scales are found, the terms in (1.7) are more conveniently determined by inserting (1.7) in (1.3) and *identifying the coefficients of* x^{-m}.

The proper choice of the asymptotic scales is essential. For instance looking for a representation of the more general form

$$\tilde{h}_a = \sum x^{-m} G_m(Ce^{-x}x^a)$$

only for $a = 1/2$ will the expansion be asymptotic (even formally) for large x. For any other value of a, $G_m(Ce^{-x}x^a) = O(x^m)$ instead of $O(1)$, and then $\tilde{h}_a$ is rather meaningless.

2 Main results

Notations. By Lemma 2 in § 3 there is a one (real) parameter family of solutions of (1.3) which are real-valued for $x > const.$ on $\mathbb{R}^+$ and such that $h_0(x) \to 0$ as $x \to +\infty$. Their behavior for large x is $h_0 \sim c_4x^{-4} + c_6x^{-6}(1 + o(1))$ where $c_4 = -392/625$ and $c_6 < 0$. We choose h_0 to be one such solution. (The choice $h_0 = \mathcal{LB}\tilde{h}_0$ has however important additional properties). We let $A \geq 9/4$ be such that, for $x > A$,

$$h_0(x) \in \left(-x^{-4}, c_4x^{-4}\right) \tag{2.1}$$

PROPOSITION 1 *(i) Any solution h of (1.3) such that $|h(x)| < 1$ for large positive x is given by*

$$h(x;C) = \sum_{k=0}^{\infty} C^k x^{-k/2} e^{-kx} h_k(x) \tag{2.2}$$

for some $C \in \mathbb{C}$, where the functions $h_k(x)$ are determined recursively as solutions of linear differential equations (cf. (3.2)) and do not depend on C.

The series (2.2) converges for all $x > x_0(C) = \max\{A, x_1\}$ where $x_1 + \frac{1}{2}\ln x_1 = \ln(|C|/12)$.
(ii) The functions h_k satisfy the inequalities

$$0 < h_k(x) < \lim_{x\to\infty} h_k(x) = \frac{k}{12^{k-1}} \quad (k \in \mathbb{N} \text{ and } x > A) \tag{2.3}$$

Furthermore,

$$\overline{\lim_{k\to\infty}} h_k(x)^{-1/k} = 12 + O(x^{-1}) \ (as\ x \to +\infty)$$

*(iii) (*Outermost pole of $h(x;C)$ on $\mathbb{R}^+$*.)*
If $C > C_0 = 12\min_{t\geq A}\sqrt{t}e^t(1-\frac{9}{4t})^{-1}$, then $h(x;C)$ has a double pole at the point $x_p = x_p(C;0)$, the unique solution of

$$x^{1/2} e^x \overline{\lim_{k\to\infty}} h_k(x)^{-1/k} = C \tag{2.4}$$

for $x > A$. The function $h(x;C)$ has no other singularities on $[x_p, \infty)$.
*(iv) (*Monotonicity and bounds for x_p*.) For $C > C_0$, $x_p(C;0)$ is increasing in C and satisfies the inequalities*

$$x_p + \frac{1}{2}\ln x_p \leq \ln(C/12) \leq x_p + \frac{1}{2}\ln x_p - \ln\left(1 - \frac{9}{4x_p}\right) \tag{2.5}$$

Thus, x_p has the asymptotic behavior,

$$x_p = \ln(C/12) - \frac{1}{2}\ln(\ln(C/12)) + O(1/\ln C) \ (C \to +\infty) \tag{2.6}$$

*(v) (*Poles of $h(x;C)$ in $\mathbb{C}$, for large C complex*.) Let $N \in \mathbb{N}$ and $\epsilon > 0$. As $|C| \to \infty$, $h(x;C)$ has double poles at the points:*

(2.7)
$$x_p(C;k) = \ln(C/12) - \frac{1}{2}\ln(\ln(C/12)) + 2k\pi i + o(1) \quad (k \in \mathbb{Z} \cap [-N, N])$$

and no poles in the region $\{x : |\Im(x)| \le N,\ \Re(x) > \Re(x_p(C;0)) + \epsilon\}$.

*(vi) (*Connection to behavior on antistokes lines.*) For the special choice* $h_0 = \mathcal{LB}\tilde{h}_0$ *we have* $h_k = \mathcal{LB}\tilde{h}_k$, *and*

$$h(x;C) \sim (C \pm \frac{1}{2}S)x^{-1/2}e^{-x} \text{ as } x \to \pm i\infty \tag{2.8}$$

where $C \in \mathbb{C}$ *is the same as in (2.2) and* S *is the Stokes constant of the* equation *(1.3).*

Notes. (1) The property that the solutions which vanish in some direction can be expressed, as in (2.2), as a power series in the constant C of (2.8) is very general [16] (integrability plays no role). Thus, for P1, C is also a natural parameter for pole location. A formula of type (2.4) is expected to be quite general and is not a connection formula stricto sensu, because it involves limits. For P1 however, there are indications that these limits can be calculated in closed form.

(2) For the choice $h_0 = \mathcal{LB}\tilde{h}_0$ it can be shown, using the methods in [17], that (2.1) holds for $x > 4/3$.

(3) The fact that P1 is integrable makes it possible to calculate explicitly the Stokes constant: $S = i\sqrt{\frac{6}{5\pi}}$. This was done in [9] using the isomonodromic deformation method; its computation using exponential is the object of a forthcoming paper. A detailed description of the asymptotic behavior of solutions of P1, derived through isomonodromy, can be found in [4, 5, 6, 9].

3 Proofs and further results

Before proving the main proposition we describe some asymptotic properties of the solutions of (1.3) which are bounded as $x \to +\infty$.

Lemma 2 *(i) If* h *is a solution of (1.3) and* $|h(x)| < 1$ *for large positive* x, *then*

$$h \sim \tilde{h}_0 = \sum_{k=4}^{\infty} c_k x^{-k} \qquad (x \to e^{i\theta}\infty \text{ with } \theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)) \tag{3.1}$$

(the first few terms of $\tilde{h}_0$ *are* $-\frac{392}{625}x^{-4} - \frac{6272}{625}x^{-6} - \frac{141196832}{390625}x^{-8} - ...$*).*
(ii) There is a one (complex/real) parameter of such solutions (complex/real valued on $\mathbb{R}^+$*); if* h_1 *and* h_2 *satisfy (3.1) then*

$$h_1 - h_2 = Ce^{-x}x^{-1/2} + o(e^{-x}x^{-1/2})$$

for some $C \in \mathbb{C}$*, as* $x \to \infty$*. If in addition* $C = 0$ *then* $h_1 = h_2$*.*

Proof of Lemma 2: (i) is standard (see, e.g. [14]); it is also easy to reprove in our case, since (1.3) can be rewritten as

$$h'' + \frac{1}{x}h' - \left(1 + \frac{1}{4x^2}\right)h = \frac{1}{2}h^2 - \frac{1}{4x^2}h + \frac{392}{625x^4} := g(x)$$

which, in integral form

$$h(x) = \frac{1}{2}x^{-1/2}\left(e^x \int_\infty^x e^{-t}t^{1/2}g(t)\mathrm{d}t - e^{-x}\int_a^x e^t t^{1/2}g(t)\mathrm{d}t\right)$$

is a contraction in the unit ball in the sup norm, for large x. If $a \in \mathbb{R}$ then h is real-valued.

Part (ii) is proven without difficulty in essentially the same way, by writing an integral contractive equation for $\delta = h_2 - h_1$ (see [16] for details under more general assumptions). ∎

3.1 Properties of (2.2) on $\mathbb{R}^+$

Let h_0 and A be as in § 2. Formal substitution of (2.2) in (1.3) yields

(3.2)
$$H_k'' - \left(1 + h_0(x) - \frac{1}{4x^2}\right)H_k = \frac{\sqrt{x}}{2}e^{-kx}\sum_{0<j<k} h_j h_{k-j} := R_k(x) \ (k \geq 1)$$

where $H_k(x) = x^{-(k-1)/2}e^{-kx}h_k(x)$.
Step 1. Study of the homogeneous part of (3.2) (same as (3.2) for $k = 1$*):*

$$y'' - \left(1 + h_0(x) - \frac{1}{4x^2}\right)y = 0 \tag{3.3}$$

Lemma 3 *(i) Eq. (3.3) has two solutions* y_+, y_- *such that:* $y_\pm(x) > 0$ *for* $x > A$*,* $y_\pm(x) = e^{\pm x}(1 + O(x^{-1}))$ *as* $x \to +\infty$ *and their Wronskian* $(y_- y_+' - y_-' y_+)$ *is* $W = 2$*.*

If $y(x)$ solves (3.3) and $y(x) = O(e^{-x})$ as $x \to \infty$ then $y = const.y_-$.
(ii) The function $u(t,s) = y_+(t)y_-(s) - y_-(t)y_+(s)$ is positive for $t > s > A$.

PROOF: (i) Except for the explicit region of positivity of $y_\pm$, this is standard asymptotic theory (see e.g. [14]; note also that $W = const$ and $W \to 2$ for $x \to \infty$). We prove the statement only for y_-; the argument for y_+ is similar. Associating to (3.3) the integral equation

$$f(x) = 1 - \int_x^\infty e^{x-s} \sinh(s-x) \left(\frac{1}{4s^2} - h_0 \right) f(s) ds \tag{3.4}$$

where $f(x) = e^x y(x)$, we define

$$J(\Psi)(x) = \int_x^\infty e^{x-s} \sinh(s-x) \left(\frac{1}{4s^2} - h_0 \right) \Psi(s) ds \tag{3.5}$$

on $L^\infty(x_0, \infty)$, for $x_0 > A$. ∎

REMARK 1 $\|J\|_{L^\infty \mapsto L^\infty} < \dfrac{1}{8x_0} \left(1 + \dfrac{1}{3x_0^2} \right)$ *and J is positivity preserving. Thus (3.4) has a unique solution f_1 in $L^\infty(A, \infty)$; moreover, $0 < f_1 < 1$ and $f_1(x) \sim 1 + \sum_{j=1}^\infty (-1)^j |c_j| x^{-j}$ $(x \to +\infty)$.*

PROOF: The first part is straightforward. Now,

$$\|f_1\|_\infty \le (1 - \|J\|)^{-1} \|1\|_\infty$$

thus $|f_1| \le \frac{6}{5}$ implying $|f_1 - 1| \le \frac{6}{5} J(1) \le \frac{1}{5}$, in particular $f_1 > 0$. Since J preserves positivity, we have $f_1 = 1 - J f_1 < 1$. The asymptotic expansion of f_1 follows from the representation $f_1 = (\sum_{j=0}^\infty (-J)^j)\, 1$. ∎

PROOF OF LEMMA 3: (ii) note that $u(t,s)$ is a solution of $u_{tt} = (1 + h_0(t) - \frac{1}{4}t^{-2})u$ with initial condition $u(s,s) = 0; u_t(s,s) = 2$. Since $1 + h_0(t) - \frac{1}{4}t^{-2} > 0$ for $t > A$, we have $u(t,s) > 0$ for all $t > s$. ∎

LEMMA 4 *There is a unique solution of (3.2) with $H_1 = y_-$ such that for $k > 0$, $H_k(x) = O(e^{-kx})$ as $x \to \infty$. Namely, the $H_k(x)$ are inductively given by*

$$H_k(x) = \frac{1}{2} \int_x^\infty u(t,x) R_k(t) \mathrm{d}t, \ k \ge 2 \tag{3.6}$$

In addition $H_k(x) > 0$ for $x > A$.

PROOF: Immediate induction using Lemma 3. Uniqueness follows from the behavior for large x of the solutions of the homogeneous equation, $y_\pm$ (Lemma 3). ∎

Step 2. Upper bounds for H_k. We have, by (3.2) and Lemma 4

(3.7)
$$H_k(x) = \int_x^\infty \sinh(t-x) R_k(t) dt - \int_x^\infty \sinh(t-x)\Big(\frac{1}{4t^2} - h_0(t)\Big) H_k(t) dt$$
$$\leq \int_x^\infty \sinh(t-x) e^{-kt} R_k(t) dt$$

From Lemma 3 and Remark 1 we get $H_1(x)e^x \in (0,1)$ on (A,∞). Assuming by induction that $H_m(x) \leq \frac{m}{12^{m-1}} x^{-(k-1)/2} e^{-mx}$ if $1 \leq m < k$, we get

$$(3.8)\quad H_k(x) \leq \frac{1}{2}\int_x^\infty \sinh(t-x) t^{-k/2+1/2} e^{-kt} \sum_{0<j<k} \frac{j(k-j)}{12^{k-1}} dt \leq$$
$$\frac{(k^3-k)}{12^k} x^{-(k-1)/2} \int_x^\infty \sinh(t-x) e^{-kt} dt = \frac{k}{12^{k-1}} x^{-(k-1)/2} e^{-kx}$$

□

Step 3. Monotonicity

LEMMA 5 *The functions $H_k(x)$ and $F_k(x) = x^{-1/2} e^{-kx} h_k(x)$, $k \geq 1$, are decreasing for $x > A$.*

PROOF: The functions F_k satisfy the equations

$$(3.9)\qquad F_k'' = (1+h_0)F_k - \frac{1}{x}F_k' + \frac{1}{2}\sum_{0<j<k} F_j F_{k-j} \qquad (k \geq 1)$$

By (3.8) and Lemma 4, the F_k are positive and vanish as $x \to \infty$. On the other hand, F_k can have no extremum for $x > A$ (otherwise, by (3.9) all extrema would be *minima*), thus are decreasing in x. For H_k the proof is very similar. ∎

Step 4. Lower bounds for H_k.

LEMMA 6 *For $x > A$ we have*

$$(3.10)\quad H_m(x) \geq \left(1-\frac{1}{8x}\right)\left(1-\frac{9}{4x}\right)^{(m-1)/2} \frac{m}{12^{m-1}} x^{-(k-1)/2} e^{-mx}$$

From Remark 1 we get $f = 1 - Jf \geq 1 - J1 \geq 1 - (1-(8x)^{-1})^{-1}$ for $x > 5/3$. Thus $H_1(x) \geq (1-(8x)^{-1})$. Since the functions H_k are decreasing, we have

$$(3.11)\quad H_k(x) \geq \int_x^\infty \mathrm{d}t \sinh(t-x) R_k(t) - H_k(x) \int_x^\infty \mathrm{d}t \sinh(t-x)\left(\frac{1}{4t^2} - h_0(t)\right)$$

Thus

$$(3.12)\quad H_k(x) \geq \left(1-\frac{1}{8x}\right) \int_x^\infty \sinh(t-x) R_k(t) \mathrm{d}t$$

From this point on, the proof of Lemma 6 follows the same track as Step 2, noting that $(1-t^{-1})^{(k-1)/2}$ is increasing in t and that $\int_x^\infty \sinh(t-x) t^{-(k-1)/2} e^{-kt} \geq (1-x^{-1})(k^2-1)^{-1} x^{-(k-1)/2} e^{-kx}$. □

Step 5 COMPLETING THE PROOF OF PROPOSITION 1 (I) AND (II): With $C \in \mathbb{C}$ given, (2.2) converges for $x > x_0(C)$, by (3.8). Furthermore, since the H_k are decreasing, if (2.2) converges at $x = x_1$, then it converges uniformly on $[x_1, \infty)$ (together with its derivatives, in view of the equality in (3.7)) and thus is a solution of (1.3). By (3.8) and (3.10), we have $h(x;C) - h_0(x) = Ce^{-x} x^{-1/2}(1+o(1))$ for large x. Therefore, by Lemma 2, if g is solution of (1.3) which vanishes as $x \to \infty$, then $g = h(x;C)$ for some $C \in \mathbb{C}$. ∎

Step 6. Divergence of the series (2.2) implies singularities of the function $h(x;C)$. Let $C > C_0$ and $x_{div} := \inf\{x | \sum_{k=1}^\infty C^k F_k(x) < \infty\}$. We have just shown that (2.2) solves (1.3) for $x > x_{div}$. By the definition of C_0 and (3.7) we have $x_{div} > A$. We now want to show that the solution $h(x;C)$ is nonanalytic at x_{div}. Let $S_N(x) = \sum_{k=1}^N C^k F_k(x)$.

LEMMA 7 *Assume $h(x;C)$ is a $C^2(x_1,\infty)$ solution of (1.3), and $x_1 \in (A, x_{div}]$.*

(i) The function $\delta_N(x) = h(x;C) - h_0 - S_N(x)$ *is positive for all* $x > x_1$ *and* $N \in \mathbb{N}^+$.

(ii) $x_1 = x_{div} = x_p(C;0)$.

PROOF: (i) We have, by Proposition 1 (ii) $\delta_N(x) = \sum_{j>N} C^j F_j(x) > 0$ for $x > x_{div}$. Now, since the function δ_N satisfies the equation

$$\delta_N'' = \frac{1}{2}\delta_N^2 + (1 + S_N + h_0)\delta_N - \frac{\delta_N'}{x} + \frac{1}{2}\sum_{\substack{i,j\in(1,N-1)\\ i+j>N}} F_i F_j \tag{3.13}$$

we see, as in Lemma 5, that δ_N is decreasing, therefore positive, for all $x > x_1$.

(ii) By (i), $h(x;C)$ is unbounded at x_{div}. But the only singularities of $h(x;C)$ for $x \neq 0$ are double poles (see the appendix). ∎

Step 7.

CONCLUDING THE PROOF OF PROPOSITION 1, (III) AND (IV):
The monotonicity of $x_p(C;0)$ in C is now immediate, while (2.5) follows from Lemma 7, (3.8) and (3.10). ∎

Note. Formula (2.4) in itself is not a practical numerical way to calculate $x_p(C)$, given a particular C. An efficient (and rigorously controllable) method is to use the procedure in this section, to obtain successively better upper and lower bounds for H_k.

3.2 Correspondence between (1.6), (2.2) and (2.8)

LEMMA 8 *If* $h_0 = \mathcal{L}\mathcal{B}\tilde{h}_0$ *then* $h_k = \mathcal{L}\mathcal{B}\tilde{h}_k$ *for all* $k \in \mathbb{N}$. *Thus the representations (1.6) and (2.2) coincide. Formula (2.8) holds.*

PROOF OF LEMMA 8 AND OF PROPOSITION 1, (VI):
We let $g_{1,2}(x,C) = h(x;C)$ as given by (1.6) and (2.2), respectively. In [16], formula (2.8) for $g_1(x,C)$ and the estimate $g_1(x,C) - h_0(x) = Ce^{-x}x^{-1/2}(1+o(1))$ are proved in a general setting. Lemma 2 then implies, as in Step 5 § 3.1, that $g_1(x;C) = g_2(x;C)$. ∎

Note. Using Lemma 8 it follows from [16] that (2.2) converges in fact for all large x with $\arg(x) \in (-\pi/2, \pi/2)$.

3.3 Uniform representation for large x

We are now interested in the region of validity of (1.7) and deriving the behavior of $x_p(C)$ for large $C \in \mathbb{C}$. While the results in the previous sections used specific properties of (1.3) (positivity of h_k), the technique of transasymptotic matching introduced in §1.2 is expected to work for more general nonlinear systems. We will only prove (1.7) to leading order, sufficient to obtain (2.7); cf. also Remark 3. The expansion (1.7) is studied in a region of x containing points near x_N (see (1.9)) and extending to infinity. Let $\epsilon > 0$ be small, $M \in \mathbb{R}^+$ and $s = s(x;C) = 12e^x x^{1/2}/C$. Define $\mathcal{R} = \{x : |\Im(x)| \le M\,|s-1| \ge \epsilon \text{ and } \Re(s) > 1-\epsilon\}$.

LEMMA 9 *Let $C_1 \in \mathbb{R}^+$ be large. For any C with $|C| > C_1$ we have*

$$h(x;C) = \frac{12s}{(s-1)^2} + \psi(x;C) \tag{3.14}$$

where $|\psi(x;C)| \le K|\ln C|^{-1}$ in Clos($\mathcal{R}$), for some K not depending on C.

Step 1: Proof of (3.14) for $x \in \mathbb{R}^+$. From (2.2) and Lemma 6 we obtain

$$h(x;C) = h_0 + \sum_{k=1}^{\infty} 12ks^{-k} + \sum_{k=1}^{\infty} 12ks^{-k}\Big[(1+r_k(x)x^{-1})^k - 1\Big] = \tag{3.15}$$

$$\frac{12s}{(s-1)^2} + \sum_{k=1}^{\lfloor x\rfloor} \frac{12k^2 c_k(x)}{x} s^{-k} + \sum_{\lfloor x\rfloor+1}^{\infty} 12k\left[(1+r_k(x)x^{-1})^k - 1\right] + O(x^{-4}) =$$

$$\frac{12s}{(s-1)^2} + \frac{E_1(x)}{\ln(C)} \quad \text{with } |E_1(x)| \le const.\ (x \in \mathbb{R}^+ \cap \mathcal{R})$$

The estimates in (3.15) are obtained in the following way: by Lemma 6 we have $|r_k(x)| \le 9/4$; thus, $|c_k(x)| = |(1+r_k/x)^k - 1| \le (e^{9/4}-1)kx^{-1}$ for $k \le x$; the tail $\sum_{\lfloor x\rfloor+1}^{\infty}(\cdot)$ is exponentially small, while for $x \in \mathcal{R}$ we have $h_0 = O(x^{-4})$ and $|x/\ln C| \le 2$ as $C \to \infty$. Similarly,

$$\left|\frac{\mathrm{d}}{\mathrm{d}x}\Big(h(x,C) - 12s(s-1)^{-2}\Big)\right| \le \frac{const}{\ln(C)} \text{ for } x \in \mathbb{R}^+ \cap \mathcal{R}) \tag{3.16}$$

Step 2. Extension to $\mathcal{R}$. Let $\psi(x) = h(x;C) - 12s/(s-1)^2$. It is convenient to change variables to $v = x + \frac{1}{2}\ln x - \ln(C/12)$ and define $\mathcal{R}_v = \{v | x \in \mathcal{R}\}$. Then $\boldsymbol{\psi} = (\psi, \psi')$ satisfies the system

$$(3.17)\quad \frac{\mathrm{d}}{\mathrm{d}v}\begin{pmatrix}\psi_1\\ \psi_2\end{pmatrix} = \begin{pmatrix}0 & 1\\ F(v) & 0\end{pmatrix}\begin{pmatrix}\psi_1\\ \psi_2\end{pmatrix} + \begin{pmatrix}0\\ \left(1+\frac{1}{2x}\right)^{-2} r\end{pmatrix} := A\boldsymbol{\psi} + \mathbf{R}$$

where $F(v) = (10e^v + e^{2v} + 1)(e^v - 1)^{-2}$ and

$$r = \frac{\psi_1^2}{2x^{1/2}} + \frac{(x+\frac{1}{4})\psi_1}{(x+\frac{1}{2})^2} + \frac{2\psi_2 - \psi_1}{4x^2} - 3\frac{e^{3v}+4e^{2v}+e^v}{(e^v-1)^4 x^{3/2}} + \frac{24e^v(2e^v+1)}{x^{1/2}(e^v-1)^4} - \frac{c_4}{x^{7/2}}$$

Let $U(t;v)$ be the fundamental matrix of $U' = AU$ (the solution with $U(t;t) = I$, the identity matrix). [2]

REMARK 2 *$U(t;v)$ and $U^{-1}(t;v)$ are uniformly bounded on*

$$\mathcal{R}_{M;t} = \{(t,v) \in \mathcal{R}_v^2 |\ |t-v| < M\}$$

PROOF: U is uniformly bounded since F is uniformly continuous on $\mathcal{R}_v$. Since $\mathrm{Tr}(A) = 0$ we have $\det(U) = const = 1$, so U^{-1} is bounded as well. ∎

Let $v_0 \in \mathcal{R}_v \cap \mathbb{R}$. By (3.15) and (3.16) we have $|\boldsymbol{\psi}(v_0)| \le const./\ln C$. We write (3.17) in integral form

$$(3.18)\qquad \boldsymbol{\psi}(v) = U(v_0;v)\boldsymbol{\psi}(v_0) + U(v_0;v)\int_{v_0}^{v} U^{-1}(v_0;t)\mathbf{R}(t)dt$$

The integral equation (3.18) is contractive in $L^\infty(\mathcal{R}_{M;v_0})$ in a ball of radius $K/\ln C$ for some K which is, by (3.15) (3.16) and Remark 2, independent of v_0 (note that all terms on the r.h.s. of (3.18) are in fact $O(1/\ln C)$ if $\|\boldsymbol{\psi}\| = 1$). Lemma 9 is proven. □

Step 3: Pole location for large $C \in \mathbb{C}$.

PROOF OF PROPOSITION 1, (V): Let

$$\mathcal{K} = \{x : |s-1| = \epsilon \text{ and } |v - 2N\pi i| < 2\pi\}$$

If ϵ is small enough, then for large C the curve $\mathcal{K}$ is a simple closed curve in $\mathrm{Clos}(\mathcal{R})$. We have by Lemma 9 $\oint_{\mathcal{K}} (\int h(\cdot;C)) = -12\pi\imath + o(1)$ thus h is not analytic in the interior of $\mathcal{K}$. The only singularities of $h(x;C)$ for $x \neq 0$ are double poles, cf. the appendix. ∎

[2] Incidentally, the equation $U' = AU$ can be solved explicitly: for the associated second order equation the general solution is $\psi = C_1 t(t+1)(t-1)^{-3} + C_2(t^{-1}P_5 + 60tv(t+1))(t-1)^{-3}$, where $t = e^v$ and P_5 is a quintic polynomial in t.

REMARK 3 *Using the asymptotic behavior of the solution of $U' = AU$ and more careful estimates, it is possible to prove contractivity and thus (3.14) in a sector of opening less than π centered on $\mathbb{R}^+$ (the opening cannot equal π because of (2.8)). The asymptotic expansion (3.14) can be determined then to higher orders in a straightforward way. (The equation for G_k is (3.17), with r.h.s. a bilinear expression in G_m, $m < k$ (cf. also footnote 2)). The result is*

$$g_C \sim G_0(s) + \frac{1}{x}G_1(s) + ... + \frac{1}{x^n}G_n(s) + ... \tag{3.19}$$

with

$$G_0 = \frac{12s}{(s-1)^2},\ G_1 = -\frac{15s^3 + 175s^2 + 30s - 2}{10s(s-1)^3},\ \cdots \tag{3.20}$$

It is also worth noting that for all $G_k(t)$ to be rational functions, the value $-392/625$ in (1.3) is crucial. A different numerical value produces a valid expansion (3.19) but with $G_k, k \geq 5$ not rational.

REMARK 4 *The arguments that were used to determine the asymptotic position of poles work even if C is not large, as long as $|x|$ is large, yielding the location of poles in the neighborhood of the imaginary axis, for large x (while on the positive real line C small implies $x_p(C)$ small, by Proposition 1).*

Appendix

All solutions of P1 are meromorphic.

The fact that the only singularities of solutions of P1 are double poles was indicated by Painlevé, but it was not until the advent of the isomonodromic approach (see [1] and [7]) that a rigorous proof was given. For completeness, we present an independent proof of this crucial property of P1 (and thus of (1.3), for $x \neq 0$), based on the idea that near poles, the equation P1 is meromorphically equivalent to an elliptic equation [18].

Notations: z_0 denotes a point where initial conditions for y are given, $\tilde{z}$ will denote singular points of y. Let $\tilde{D}$ be a disk without a discrete set of points. $\Lambda(z_1, z_2)$ will denote smooth curves in $\tilde{D}$, with the endpoints z_1 and $z_2 \in \mathrm{Clos}(\tilde{D})$ removed, such that the length of any subsegment $\Lambda(z_3, z_4)$ of $\Lambda(z_1, z_2)$ is less than $2|z_3 - z_4|$.

LEMMA 10 *In a neighborhood of any point $\tilde{z} \in \mathbb{C}$ there exists a one-parameter family of solutions h of (1.1), having a double pole at $\tilde{z}$, of the form $h(z) = (z-\tilde{z})^{-2}A(z)$ where $A(z)$ is analytic at $\tilde{z}$, and $A(\tilde{z}) = 1$.*

PROOF: Substituting $y = \sum_{k=-2}^{\infty} c_k (z-\tilde{z})^k$ in (1.1) we get

$$(k+1)(k+2)c_{k+2} = 6\sum_{m=-2}^{k+2} c_m c_{k-m} + \tilde{z}\delta_{k=0} + \delta_{k=1} \tag{A.1}$$

so that $c_{-2} = 0$ or 1; choosing $c_{-2} = 1$ we have $c_{-1} = 0$ and

$$[(k+1)(k+2) - 12]c_{k+2} = 6\sum_{m=0}^{k} c_m c_{k-m} + \tilde{z}\delta_{k=0} + \delta_{k=1} \tag{A.2}$$

Carrying on the calculation we get

$$\Big(c_{-2}, c_{-1}, c_0, c_1, c_2, c_3, c_4\Big) = \Big(1, 0, 0, 0, \frac{-1}{10}\tilde{z}, -\frac{1}{6}, c_4\Big) \tag{A.3}$$

i.e., c_4 is a free constant (*Note*: The specific form of the equation (1.1) is crucial in obtaining an integer power series solution. Generic perturbations of (1.1) lead to an inconsistent equation for c_4.

Finally, for some $\rho > 0$ and all $k \geq 0$ we have $|c_k| \leq \frac{1}{3}(k+1)\rho^{k+2}$. Indeed, let ρ be such that $|c_i| \leq \frac{1}{3}(i+1)\rho^{i+2}$ for $i \leq 2$. Then, assuming this property for $c_0, c_1, ..., c_{k+1}$ we have

$$|c_{k+2}| \leq \frac{1}{3}\frac{(k+1)(k+2)(k+3)}{(k-2)(k+5)}\rho^{k+4} \leq (k+3)\rho^{k+4}$$

∎

LEMMA 11 *If y is a solution of (1.1), analytic at z_0, then the radius of analyticity at z_0 is at least $R = 1/\max\{|y(z_0)|^{1/2}, |y'(z_0)/2|^{1/3}, |y_0^2 + z_0/6|^{1/4}\}$.*

PROOF: With $y = \sum_{k=0}^{\infty} c_k (z-z_0)^k$ we have

$$(k+1)(k+2)c_{k+2} = 6\sum_{m=0}^{k} c_m c_{k-m} + z_0\delta_{k=0} + \delta_{k=1} \tag{A.4}$$

By the definition of R we have $|c_j| \leq (j+2)R^{-j-2}$ for $j \leq 2$. Inductively, if the inequality holds $\forall j \leq k+1$ we have

$$|c_{k+2}| \leq \frac{(k+1)(k+2)(k+3)R^{-k-4}}{(k+1)(k+2)} = (k+3)R^{-k-4}$$

∎

LEMMA 12 *Let y be a solution of (1.1), analytic at any $z \in \Lambda(z_0, \tilde{z})$ and that $y(z) = \sum_{k=-2}^{4} a_k(z-\tilde{z})^k + O(z-\tilde{z})^5$ as $z \to \tilde{z}$ along $\Lambda(z_0, \tilde{z})$, with $a_{-2} \neq 0$.*

Then y is meromorphic at $\tilde{z}$, and y is uniquely determined by $\tilde{z}$ and a_4.

PROOF: Since $\sum_{k=-2}^{4} a_k(z-\tilde{z})^k$ is an $O(z-\tilde{z})^5$ formal solution of P1, for $i \leq 3$ we must have $a_k = c_k$ defined in (A.3) (note: since y solves (1.1), the asymptotic expansion is differentiable). Let $y_0(z)$ be the solution given in Lemma 10 with $c_4 = a_4$. Let $v(z) = y_0(z) - (z-\tilde{z})^{-2}$ and $f(z) = y(z) - y_0(z)$. Then $v = O((z-\tilde{z})^2), f = O(z-\tilde{z})^5$ and

$$f'' - \frac{12}{(z-\tilde{z})^2} f = 12fv + 6f^2 \tag{A.5}$$

or

$$f(z) = C_1(z-\tilde{z})^{-3} + C_2(z-\tilde{z})^4 - \frac{7}{(z-\tilde{z})^3}\int_{\tilde{z}}^{z} (s-\tilde{z})^4 g(s)ds$$
$$+\frac{(z-\tilde{z})^4}{7}\int_{\tilde{z}}^{z} (s-\tilde{z})^{-3} g(s)ds = \mathcal{N}(f) \tag{A.6}$$

where $g = 12fv + 6f^2 = O(z-\tilde{z})^7$ near $\tilde{z}$. Thus $C_1 = C_2 = 0$. Let $z_1 \in \Lambda$ and $\mathcal{S}$ be the Banach space of continuous functions on $\Lambda(z_1, \tilde{z})$ with the norm $\sup_\Lambda |(z-\tilde{z})^{-5} f(z)|$. It is immediate to check that if $z_1 - \tilde{z}$ is small enough $\mathcal{N}$ is contractive in the unit ball $B \subset \mathcal{S}$. The solution of (A.6) is thus unique in B, $f = 0$. ∎

Let y be a solution of (1.1), and D an open disk where y is meromorphic. By definition, the singularities of y in D are isolated. Let $\tilde{z} \in \partial D$ be a singular point of y. We will show that $\tilde{z}$ is a pole: *this implies that y is meromorphic on $\mathbb{C}$.*

Let $\tilde{D}$ be D without the poles of y.

REMARK 5 *(i) As $z \to \tilde{z}$ along $\Lambda(z, \tilde{z})$ we have $|y(z)| + |y'(z)| \to \infty$. (ii) y is unbounded as $z \to \tilde{z}$.*

PROOF: (i) Since y is singular at $\tilde{z}$ then by Lemma 11 $y(z)$ or $y'(z)$ must be large for z near $\tilde{z}$. (ii) Multiplication of (1.1) by y' and integration yields:

$$(y')^2 = 4y^3 + 2zy - 2\int_a^z y(s)ds + C \tag{A.7}$$

If y were bounded along Λ, then by (A.7) y' would be bounded as well. ∎

PROPOSITION 13 *Let $z_0 \in D$ be a regular point of y and $\Lambda = \Lambda(z_0, \tilde{z}) \subset D$. We have $y^{-3}(s)\int_a^s y(t)dt \to 0$, $|y(s)| \to \infty$, and $|y'(s)| \to \infty$ as $s \to \tilde{z}$ on Λ.*

PROOF: For *any* continuous function f unbounded at $\tilde{z}$ we have

$$\liminf_{z\to\tilde{z}} |f|^{-3}\int_a^z |f|(t)d|t| \le \liminf_{z\to\tilde{z}} |f|^{-3} \max_{\Lambda(a,z)} |f| \,\text{length}(\Lambda) = 0$$

Thus, if we had $y^{-3}(s)\int_a^s y(t)dt \not\to 0$, then for any $\gamma > 0$ small enough

$$\left|\int_a^{s_n} y(t)dt\right| = \gamma |y^3(s_n)| \tag{A.8}$$

for some sequence $s_n \to \tilde{z}$ on Λ. If $\gamma < \frac{1}{2}$, by (A.8), (A.7) and Remark 5 we have

$$|y(s_n)| \to \infty \ \text{ and } \ |y'(s_n)| \to \infty \ \text{ and } \ \int_{z_0}^{s_n} |y(t)|d|t| \to \infty \tag{A.9}$$

Consider now the solution h of the equation

$$(h')^2 = 4h^3 + 2s_n h + \tilde{C} \tag{A.10}$$

satisfying $h(0) = y(s_n), h'(0) = y'(s_n)$ (with $\tilde{C}$ chosen accordingly in (A.10)). It follows from (A.9) that $\tilde{C} = C - 2\int_a^{s_n} y(s)ds \to \infty$. Define the function H by

$$\alpha H(\beta z) := h(z), \ \text{ where } \alpha = \tilde{C}^{1/3}, \ \beta = \tilde{C}^{1/6}$$

It follows that

$$(H')^2 = 4H^3 - 2\theta H + 1, \text{ where } \theta = \tilde{C}^{-2/3} \tag{A.11}$$

H is thus an elliptic function with with periods $\Pi_{1,2} = \Pi + o(\theta)$ for some $\Pi \in \mathbb{C}$ (independent of θ). By (A.8) $|y(s_n)| = (2\gamma^{-1/3})|\tilde{C}^{1/3}|(1 + o(1))$, then using (A.7), we get $|y'(s_n)| = (4\gamma^{-1}e^{i\phi_n} + 2)^{-1/2}|\tilde{C}^{1/2}|(1 + o(1))$ for some $\phi_n \in \mathbb{R}$.

Therefore $|H(0)| = (2\gamma)^{-1/3} + o(1)$, and $|H'(0)| = (4\gamma^{-1}e^{i\phi_n} + 2)^{-1/2}| + o(1)$. With the substitution $\alpha Y(\beta z) := y(s_n + z)$ we have

$$(Y')^2 = 4Y^3 + 2\theta Y + 1 + 2\epsilon \left(zY - \int_0^z Y(t)dt \right), \quad \text{where } \epsilon = \tilde{C}^{-5/6},$$
$$Y'' = 6Y^2 + \theta + \epsilon s \tag{A.12}$$

We substitute $Y = H \circ G$; $G(z) = z + g(z)$ in (A.12) with $g(0) = g'(0) = 0$ and

$$\begin{aligned} &2g'H'' \circ G + (g')^2 H'' \circ G + g''H' \circ G = \epsilon z \\ &\text{or } 2g'(H' \circ G)' + g''H' \circ G = \epsilon z + (g')^2 H'' \circ G \end{aligned} \tag{A.13}$$

which can be rewritten in the integral form ($g(z) = \int_{\Lambda(0,z)} w(t)dt, G = z + g$):

$$w = (H' \circ G)^{-2} \int_{\Lambda(0,z)} H'(G(u)) \left[\epsilon u + H''(G(u))w^2(u)^2 \right] du = \mathcal{N}(w) \tag{A.14}$$

Let Δ_ϵ be the disk centered at 0, of diameter $|\epsilon|^{-1/4}$, and $\tilde{\Delta}_\epsilon \ni 0$ be Δ_ϵ without the disks of diameter $d(\gamma)$ around the poles and zeros of $H'(z)$. Let $\Lambda \subset \Delta_\epsilon$, let $a \in (1/4, 1/2)$. Let $\mathcal{S}$ be the Banach space of continuous functions on Λ in the sup norm and $B = \{w : \|w\| \le |\epsilon|^a\} \subset \mathcal{S}$. If $w \in B$, then $|g| \le |\int_{\Lambda(0,z)} w(t)dt| \le |\epsilon|^{a-1/4}$. B is preserved by $\mathcal{N}$ provided that

$$|\epsilon|\, Const(\gamma)|\epsilon|^{-1/2} + Const(\gamma)|\epsilon|^{2a-1/4} \le |\epsilon|^a \tag{A.15}$$

which holds for small ϵ. Similarly for $u, v \in B$, $\|\mathcal{N}(u) - \mathcal{N}(v)\| = O(\epsilon^{a-1/4})\|u - v\|$. $\mathcal{N}$ has thus a unique fixed point in B.

Therefore $|g(z)| \le |\epsilon|^{a-1/4}$ for $z \in \tilde{\Delta}_\epsilon$ and thus (when ϵ is small)

(A.16)
$$\sup_{\tilde{\Delta}_\epsilon} |\alpha^{-1} y(s_n + \beta^{-1}\cdot) - H(\cdot)| \;=\; \sup_{\tilde{\Delta}_\epsilon} |H(G(\cdot)) - H(\cdot)| \;=\; O(\epsilon^{a-1/4})$$

Note now that as $s_n \to \tilde{z}$ we have $\beta^{-1} = \tilde{C}^{-1/6} \to 0$ while $\alpha \to \infty$ and $|\beta^{-1}\epsilon^{-1/4}| \propto \tilde{C}^{1/24} \to \infty$. Thus $L = s_n + \alpha^{-1}\tilde{\Delta}_\epsilon$ is an $\alpha^{-1}-$lattice for D (i.e., $\forall z \in D\,, \exists z_1 \in L : |z - z_1| \leq \alpha^{-1}$), and (A.16) implies y is unbounded near z_0, contradiction.

Since $y^{-3}(s)\int_a^s y(t)dt \to 0$, the fact that $\min\{|y|, |y'|\} \to \infty$ now follows from Remark 5 and (A.7). ∎

PROPOSITION 14 *$\tilde{z}$ is a second order pole.*

PROOF: The substitution $y = 1/v^2$ transforms (A.7) into

$$(v')^2 = 1 + \frac{1}{2}v(z)^4 z - \frac{1}{2}v(z)^6 \int_{z_0}^{z} v(s)^{-2} ds + Cv(z)^6 \tag{A.17}$$

It follows from Proposition 13 that along a curve $\Lambda(z_0, \tilde{z})$ avoiding the poles of y we have $v^6 \int_{z_0}^{z} v^{-2}(s)ds \to 0$. With $\zeta = z - \tilde{z}$, (A.17) implies $(v')^2 = 1 + o(1)$ so that $v^2 = \zeta^2(1 + o(1))^2$ near $\tilde{z}$. Using (A.17) again we now get $v^2 = (\zeta + \sum_{k=2}^{6} v_k \zeta^k + o(\zeta^6))$. Lemma 11 implies that y is meromorphic at $\tilde{z}$. ∎

Acknowledgment.

The author would like to thank Professors Percy Deift and Martin Kruskal for very interesting discussions on this subject.

Bibliography

[1] Fokas, A. S., Mugan, U., and Zhou, X. On the Solvability of PI, III, and V *Inverse Problems* 8: 757–785, 1991

[2] Jimbo, M. and Miwa, T. Monodromy Preserving Deformation of Linear Ordinary Differential Equations with Rational Coefficients I, I *Physica D* 2: 306–352, 1981

[3] Jimbo, M. and Miwa, T. Monodromy Preserving Deformation of Linear Ordinary Differential Equations with Rational Coefficients II, I *Physica D* 2: 407–448, 1981

[4] Kapaev, A. A. and Kitaev, A. V., Connection Formulae for the First Painlevé Transcendent in the Complex Domain *Lectures in Mathematical Physics* 27: 243–252, 1993

[5] Kitaev, A. V. The method of Isomonodromy Deformations and the Asymptotics of Solutions of the "Complete" Thirf Painlevé Equation *Math. USSR Sbornik* 62: 421–444, 1989

[6] Kitaev, A. V. Elliptic Asymptotics of the First and Second Painlevé Transcendents *Russian Math. Surveys* 49:1: 81–150, 1994

[7] Its, A. and Novokshenov, V. Y. *The isomonodromic deformation method in the theory of Painlevé equations.* Springer-Verlag, 1986

[8] Écalle, J. *Fonctions Resurgentes.* Publications Mathematiques D'Orsay 81, 1981

[9] Kapaev, A. A. Asymptotics of Solutions of the Painlevé Equation of the First Kind. *Diff. Equ.* 24: 1107–1115, 1989 (Translation from the Russian original (1988).)

[10] Flaschka, H. and Newell, A. C. Monodromy- and spectrum-preserving deformations. I *Comm. Math. Phys.* 76: 65–116, 1980

[11] Écalle, J. *Finitude des cycles-limites et accéléro-sommation de l'application de retour. Lecture Notes in Math.* **1455**, Springer Berlin, 1990

[12] Écalle, J. in *Bifurcations and periodic orbits of vector fields NATO ASI Series, Vol. 408* pp. 75–184, Kluwer Academic Publishers Dordrecht; Boston, 1993

[13] Écalle, J. and Menous, F. in *The Stokes phenomenon and Hilbert's 16th problem.* pp. 71–101, World Sci. Publishing River Edge, NJ, 1995

[14] Wasow, W. *Asymptotic expansions for ordinary differential equations.* Interscience Publishers New York, 1968

[15] Costin, O. Exponential asymptotics, trans-series and generalized Borel summation for analytic nonlinear rank one systems of ODE's. *Internat. Math. Res. Notices* 8: 377–417, 1995[3]

[16] Costin, O. On Borel summation and Stokes phenomena for rank one nonlinear systems of ODE's. *Duke Math. J., to appear* [3]

[17] Costin, O. and Kruskal, M. D. On optimal truncation of divergent series solutions of nonlinear differential systems, *submitted to Proc. Roy. Soc. Lond.*[3]

[18] O. Costin , R. D. Costin Singular normal forms of nonlinear equations, *submitted to Nonlinearity*[3].

Received Month 199X.

[3]Preprints available at http://www.math.uchicago.edu/∼costin