

**Quiz 6**

**Instructions:** This quiz is worth 35 points and the value of each question is listed with each question. You may use any notes or books but you must work individually. The only computation aid which you may use is MATLAB, unless otherwise indicated. Make sure to write clearly and justify your answers.

(1.)(7 pts.) Let  $S$  be the subspace of  $\mathbb{R}^5$  spanned by  $\mathbf{v}_1 = (1, -1, -1, 2, 3)^T$ ,  $\mathbf{v}_2 = (2, 1, 0, 1, -1)^T$  and  $\mathbf{v}_3 = (-1, 5, 3, 0, 3)^T$ .

- (a.) Verify that  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is an orthogonal basis for  $S$ .
- (b.) Find the orthogonal projection  $\mathbf{p}$  of  $\mathbf{u} = (1, 2, 1, 2, 1)^T$  onto  $S$ .
- (c.) Find the norm of the residual vector  $\mathbf{u} - \mathbf{p}$  where  $\mathbf{u}$  and  $\mathbf{p}$  are as in (b.).

(2.)(10 pts.) Let  $V$  be the vector space  $C[0, 2]$  with the usual inner product and let  $S$  be the subspace of  $V$  spanned by  $\mathbf{v}_1 = 1$ ,  $\mathbf{v}_2 = x - 1$  and  $\mathbf{v}_3 = 3x^2 - 6x + 2$

- (a.) Verify that  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is an orthogonal basis for  $S$ .
- (b.) Find the orthogonal projection  $\mathbf{p}$  of  $\mathbf{u} = \sqrt{x}$  onto  $S$ .
- (c.) Find the norm of the residual vector  $\mathbf{u} - \mathbf{p}$  where  $\mathbf{u}$  and  $\mathbf{p}$  are as in (b.).

(3.)(10 pts.) Let  $f(x) = |x|$ . Find the  $n$ th order Fourier series approximation of  $f(x)$ .

(4.)(8 pts.) Suppose that a system is modeled by a function of the form  $f(x) = a + b \sin(x) + c \cos(x)$ . The following four data points  $(x, f(x))$  are collected:  $(0, -12)$ ,  $(\frac{\pi}{4}, 2)$ ,  $(\frac{7\pi}{12}, \sqrt{6})$  and  $(\frac{5\pi}{6}, 2\sqrt{3})$ .

- (a.) Find the best approximation of  $f(x)$  using the method of least squares.
- (b.) Find the approximate value of  $f(2)$  using the result from part (a.)