

Review 3

- (1.) Let $\mathbf{x} = (3, 1, -1, 5)^T$ and $\mathbf{y} = (2, 6, -4, 5)^T$ be vectors in \mathbb{R}^4 with the usual inner product. Find:
- (a.) $\langle \mathbf{x}, \mathbf{y} \rangle$, $\|\mathbf{x}\|$, $\|\mathbf{y}\|$,
 - (b.) the orthogonal projection of \mathbf{x} onto the line spanned by \mathbf{y} and
 - (c.) the angle between \mathbf{x} and \mathbf{y} .

- (2.) Let $\mathbf{f} = x$ and $\mathbf{g} = \ln(x)$ be vectors in $C[1, e]$ with the usual inner product. Find:
- (a.) $\langle \mathbf{f}, \mathbf{g} \rangle$, $\|\mathbf{f}\|$, $\|\mathbf{g}\|$,
 - (b.) the orthogonal projection of \mathbf{f} onto the line spanned by \mathbf{g} and
 - (c.) the angle between \mathbf{f} and \mathbf{g} .

- (3.) Let $A = \begin{pmatrix} 1 & 2 & 2 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{pmatrix}$, $B = \begin{pmatrix} -4 & 1 & 1 \\ -3 & 3 & 2 \\ 1 & -2 & -2 \end{pmatrix}$ and $C = \begin{pmatrix} 1 & 1 & -1 \\ 2 & 2 & 2 \\ -2 & 0 & -1 \end{pmatrix}$ be vectors in $\mathbb{R}^{3 \times 3}$ with the usual inner product.

- (a.) Verify that $\{A, B, C\}$ form an orthogonal basis for $S = \text{Span}(A, B, C)$ and
 - (b.) find the orthogonal projection of $D = \begin{pmatrix} 1 & -2 & 3 \\ 2 & 4 & 5 \\ 0 & 1 & -2 \end{pmatrix}$ onto S .
 - (c.) Compute the relative residual for the projection in part (b.)
- (4.) Let $\mathbf{f}_1 = 1$, $\mathbf{f}_2 = 2x - 1$ and $\mathbf{f}_3 = 12x^2 - 12x + 2$ be vectors in $C[0, 1]$ with the usual inner product.
- (a.) Verify that $\{\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3\}$ form an orthogonal basis for $S = \text{Span}(\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3)$ and
 - (b.) find the orthogonal projection of x^3 onto S .
 - (c.) Compute the relative residual for the projection in part (b.)

- (5.) Suppose that $f(x) = \begin{cases} -1 & -\pi \leq x < 0 \\ 1 & 0 \leq x \leq \pi \end{cases}$. Find the n th order Fourier series for $f(x)$ over $[-\pi, \pi]$.

- (6.) To measure the takeoff performance of an airplane, the horizontal position of the plane was measured every second from $t = 0$ to $t = 5$. The positions in meters were: 0, 29.9, 104.7, 222.0, 380.4 and 571.7
- (a.) Find the least squares cubic curve $y = \beta_0 + \beta_1 t + \beta_2 t^2 + \beta_3 t^3$ for these data.
 - (b.) Use the result of (a.) to estimate the velocity of the plane when $t = 4.5$ seconds.

- (7.) Let $A = \begin{pmatrix} 7 & -6 & -4 & 1 \\ -5 & 1 & 0 & -2 \\ 10 & 11 & 7 & -3 \\ 19 & 9 & 7 & 1 \end{pmatrix}$

- (a.) Find the condition number of A with respect to the 1-norm on A .
 - (b.) Find the condition number of A with respect to the ∞ -norm on A .
 - (c.) Is this matrix well-conditioned or ill-conditioned? Explain your answer.
- (8.) Prove:
- (a.) If S is subspace of the inner product space V , then the orthogonal complement S^\perp of S is also a subspace.
 - (b.) If \mathbf{x} and \mathbf{y} are orthogonal vectors in the inner product space V , then $\|\mathbf{x} - \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2$.
 - (c.) If \mathbf{p} is the projection of \mathbf{x} onto the line spanned by \mathbf{y} , then \mathbf{p} is orthogonal to $\mathbf{x} - \mathbf{p}$.

(9.) True or False:

- (a.) If A is an $m \times n$ matrix, then $\dim(\text{Col}(A)) + \dim(\text{N}(A^T)) = n$.
- (b.) If \mathbf{p} is the orthogonal projection of \mathbf{x} onto the subspace S , then \mathbf{p} and $\mathbf{p} + \mathbf{x}$ are orthogonal.
- (c.) The matrix equation $A\mathbf{x} = \mathbf{b}$ has a unique least squares solution $\hat{\mathbf{x}}$.
- (d.) If V is a normed linear space and \mathbf{x} and \mathbf{y} are vectors in V , then $\|\mathbf{x} + \mathbf{y}\| \geq \|\mathbf{x}\| + \|\mathbf{y}\|$.
- (e.) If V is an inner product space and S is a finite dimensional subspace of V , then S has an orthonormal basis.

(10.) Suppose that $A = \begin{pmatrix} -6 & 3 & -27 & -33 & -13 \\ 6 & -5 & 25 & 28 & 14 \\ 8 & -6 & 34 & 38 & 18 \\ 12 & -10 & 50 & 41 & 23 \\ 14 & -21 & 49 & 29 & 33 \end{pmatrix}$.

- (a.) Find a basis for $\text{Col}(A)$.
- (b.) Find a basis for $\text{Col}(A)^\perp$.